PINEHURST MEDICAL CLINIC ENDOSCOPY CENTER PHONE: 910-295-9207

Appointment Date:	Arrival Time:
Landan 45 Dadaral Dr. Bindanat NC	Dlandalan.
Location: <u>15 Regional Dr. Pinehurst NC</u>	Physician:
Patient Name:	

GATORADE SPLIT DOSING PREP INSTRUCTIONS FOR COLONOSCOPY

Please read the following instructions carefully.

Special Medication Instructions: Continue taking ASPIRIN as directed.

If you take any blood thinners such as **Coumadin**, **Plavix**, **Effient**, **Eliquis**, **Xarelto**, **Brilinta**, **Aggrenox**, **Arixtra**, **Sayvays**a or **Pradaxa** and you did not see one of our GI providers in clinic, please contact our office immediately. You will need special instructions concerning these medications. **HOLD_____ DAYS**.

<u>GLP-1 Medications</u>: If you take any GLP-1 medications such as **Byetta, Victoza, Saxenda, Bydureon, Trulicity, Ozempic, Wegovy, Mounjaro, Zepbound, Adlyxin** you will need to hold these medications <u>10 days</u> prior to your procedure appointment. If you have any questions, please contact our office, and ask to speak with an Endoscopy Nurse.

If you take a dual insulin/GLP-1 such as **Soliqua**, please **contact your PCP** for insulin coverage during these 10 days.

Diabetic Patients: DO NOT TAKE oral diabetic medications or short acting insulin the day before your procedure. If you normally take long acting insulin in the evening, check your blood sugar. If your blood sugar is above 100, take ½ your normal dose of long acting insulin. **DO NOT TAKE** any diabetic medications the morning of your procedure. Please contact our office and speak with a nurse if you have questions concerning your diabetic medications during this period of time.

One Week Prior to Your Colonoscopy

Stop taking **Iron tablets**. This does not include multivitamins.

Three Days Prior to Your Colonoscopy

You will need to **avoid** fruits and vegetables with seeds and skins such as tomatoes and grapes, roughage foods such as leafy green vegetables or salads, and all beans, corn, and nuts. You **may** eat MEAT, WHITE BREAD, PASTA, RICE, and POTATOES.

GATORADE PREP INSTRUCTIONS

*Please Note: You will need to purchase the following (no prescriptions are needed)

- Two 32-ounce bottles of Gatorade (not red or purple)
- Diabetics should buy sugar-free Gatorade
- Two 10-ounce bottles of Magnesium Citrate Saline Laxative Oral Solution
- Dulcolax 5mg laxative tablets
- Miralax 8.3-ounce powder laxative

The Day Before the Colonoscopy

Drink only clear liquids for breakfast, lunch, and dinner. No solid foods allowed. Avoid dairy products or anything red or purple.

Clear liquid list

Water or seltzer, tea or black coffee **NO** milk or non-dairy creamer. You may have sugar or sugar substitute. Soft Drinks -Regular or diet sodas (orange, ginger ale, cola, Sprite, 7-Up, etc)
Sports drinks or clear-colored fruit drinks without pulp - Gatorade, Crystal Light, apple juice, etc.

Broths - chicken, beef, or vegetable (without any solid food such as meat, noodles, rice, or vegetables) Desserts - Jell-O (lemon, lime, orange; no fruit or toppings), popsicles, Italian ice

Preparing the Gatorade Prep Solution: In a large pitcher mix two bottles of Gatorade and 8.3oz of Miralax. Stir/shake until Miralax is completely dissolved. Chill in refrigerator.

FIRST DOSE: The Night Before Your Procedure

At 5pm: Take 3 Dulcolax 5mg laxative tablets and drink one bottle of Magnesium Citrate **At 6pm:** Drink one 8-oz glass of Gatorade prep solution every 15 minutes until half of the pitcher is gone.

SECOND DOSE: The Morning of Your Procedure_

At least five hours before your procedure drink one bottle of Magnesium Citrate followed by the remaining Gatorade prep solution; one (8 oz) glass every 15 minutes until all the Gatorade prep is gone. Make sure you drink additional liquids to ensure proper hydration. You may continue drinking clear liquids **up to 3 hours before** your procedure, then nothing by mouth.

You May Take the Following Medications with a small sip of water before your procedure: Blood Pressure, Heart, Asthma, Breathing, Seizure, Anti-Depressants, Thyroid Medications, and Aspirin.

ADDITIONAL INSTRUCTIONS:

- 1. Bring a list of your medications with the dosages to your procedure appointment.
- 2. Please bring a Photo ID and your insurance card(s).
- 3. Wear comfortable clothes that are easy to get into and out of. You may want to wear or bring socks.
- 4. Your driver may want to bring a drink and/ or snacks. We do not have snack machines in the waiting room.
- 5. Do not wear necklaces, chains, bracelets, or earrings. You may wear your wedding band and a watch.
- 6. Please remove nose and/or tongue piercings prior to your procedure.

YOU MUST HAVE A RESPONSIBLE DRIVER TO ACCOMPANY YOU!

The Driver must be **18 years or older** and stay in the waiting area. Patients cannot be dropped off and picked up later. Procedures will be delayed until the designated driver is in the waiting room.

What To Expect Prior To Your Appointment

CONFIRMATION OF APPOINTMENT:

- 1. You will receive an automated telephone reminder two weeks prior to the scheduled date of your procedure. During this phone call you will be able to confirm your appointment. If you do not get this call you must call us at (910) 235-3177 to confirm your appointment.
- 2. You must confirm your appointment no later than 7 days prior to your procedure date. If you do not, your procedure will be cancelled.
- 3. If you confirm your appointment and then fail to arrive for your procedure and you have not given us at least 72 hour notice you will be charged \$100.00 for not keeping your procedure appointment.
- 4. Please review these instructions carefully and write down any questions you may have. It is very important that you begin to follow these instructions at least 10 days prior to your appointment to ensure that you understand it and are able to follow the directions. We may be reached at (910) 235-3177 if you have additional questions.

TWO BUSINESS DAYS BEFORE YOUR PROCEDURE:

You will receive a phone call from one of our nurses to remind you of your arrival time.

NOTE: This is an elective procedure. Patients must be aware that delayed complications may occur after the procedure, especially if polyps are removed or biopsies are taken. The risk of complications is very low however your doctor recommends avoiding travel for 14 days after your procedure